Subject Index for *Journal of Ichthyology*, Volume 35, 1995

Abramis brama: **2**, 111–119; **3**, 101–108; **8**, 1–13; **9**, 69–78
Acclimation temperature: **6**, 108–118
Acclimatization: **7**, 122–142
Accumulation: **9**, 154–170
Acidified water effect: **3**, 137–139
Acipenser gueldenstaedti: **6**, 100–107; **8**, 123–136
Acipenser ruthenus: **3**, 109–120; **9**, 8–19
Acipenser stellatus: **1**, 52–58; **9**, 281–294
Acipenseridae: **6**, 72–87; **8**, 263–270
Adrenergic regulation: **8**, 1–13
Age 0+ carp: **6**, 108–118
Age-size composition; **6**, 20-27
Age-specific mortality dynamics: **9**, 249-255
Agonidae: **8**, 245–247
Alburnus alburnus: **8**, 173–183
Alepocephalidae: **6**, 130–139
Alpine char: **7**, 1–9
Amazonian rivers: **9**, 227–248
American plaice: **5**, 40–54
Amitriptyline: **9**, 392–394
Amur River basin: **9**, 374–378
Anadromous fishes: **5**, 13–27
Anadyr–Navarin region: **4**, 9–17
Analysis of possible formation tracks: **8**, 33–43
Anchovy aggregations: **3**, 151–156
Annotated list: **1**, 81–123
Annual changes: **1**, 32–43
Anomalies in fishes: **5**, 147–158
Anoplopoma fimbria: **3**, 146–150
Antarctic dragonfishes phylogeny: **5**, 89–104
Antarctic dragonfishes systematics: **5**, 89–104
Antidepressant effect: **9**, 392–394
Aphanopinae: **9**, 183–192
Arctic char: **7**, 1–9
Arctic grayling: **9**, 220–226
Arctic seas: **1**, 81–123
Arcto-Norwegian cod: **7**, 62–74
Arctozenus rissoi: **4**, 88–96; **6**, 1–19
Artedidraco mirus: **9**, 395–399
Astronesthes chrysophekadion: **2**, 21–39
Astronesthidae: **3**, 121–123
Atlantic halibut: **4**, 150–152
Atlantic salmon: **7**, 52–61

Baikal oilfishes: **1**, 139–145
Baikal omul: **3**, 54–64
Banded flounder biology: **3**, 22–28
Barbourisia rufa: **6**, 140–150
Barbourisiidae: **6**, 140–150
Barents Sea: **4**, 150–152; **5**, 40–54; **6**, 28–44, 45–60; **8**, 245–247
Bathyal region: **4**, 130–134, 135–139
Bathydraconidae: **5**, 89–104
Bathymetric distribution: **9**; 295–304
Behavior of fishes: **1**, 124–138
Behavior of roach: **4**, 121–129
Behavior studies: **2**, 75–98
Beluga sturgeon status: **9**, 59–68
Benthopelagic sculpins: **3**, 54–64
Bering Sea: **3**, 65–77; **4**, 9–17, 18–27; **8**, 24–32
Biochemical estimate: **5**, 136–146
Biological characterization: **4**, 28–45
Biological markers: **1**, 44–51
Biology of cisco: **9**, 368–373
Biology: **6**, 1–19; **8**, 44–52
Biomass: **8**, 44–52
Black Sea: **4**, 1–8, 107–120
Black Sea areas: **9**, 113–122
Bleak: **8**, 173–183
Blicca bjoerkna: **9**, 356–361
Blood serum: **2**, 1–7; **6**, 100–107; **8**, 263–270
Blunt-snouted lenok: **3**, 1–21
Bol'shaya River estuary: **2**, 65–74; **3**, 22–28; **6**, 88–99; **9**, 102–112;
Bottom fishes: **9**, 205–219
Bream: **8**, 1–13
Broken Spur hydrothermal vent field: **9**, 328–332
Bureya River: **9**, 374–378

C. (*Cichlasoma*) *portalegrense*: **7**, 86–107

Subject Index for *Journal of Ichthyology*, Volume 35, 1995

Abramis brama: **2**, 111–119; **3**, 101–108; **8**, 1–13; **9**, 69–78
Acclimation temperature: **6**, 108–118
Acclimatization: **7**, 122–142
Accumulation: **9**, 154–170
Acidified water effect: **3**, 137–139
Acipenser gueldenstaedti: **6**, 100–107; **8**, 123–136
Acipenser ruthenus: **3**, 109–120; **9**, 8–19
Acipenser stellatus: **1**, 52–58; **9**, 281–294
Acipenseridae: **6**, 72–87; **8**, 263–270
Adrenergic regulation: **8**, 1–13
Age 0+ carp: **6**, 108–118
Age-size composition; **6**, 20-27
Age-specific mortality dynamics: **9**, 249-255
Agonidae: **8**, 245–247
Alburnus alburnus: **8**, 173–183
Alepocephalidae: **6**, 130–139
Alpine char: **7**, 1–9
Amazonian rivers: **9**, 227–248
American plaice: **5**, 40–54
Amitriptyline: **9**, 392–394
Amur River basin: **9**, 374–378
Anadromous fishes: **5**, 13–27
Anadyr–Navarin region: **4**, 9–17
Analysis of possible formation tracks: **8**, 33–43
Anchovy aggregations: **3**, 151–156
Annotated list: **1**, 81–123
Annual changes: **1**, 32–43
Anomalies in fishes: **5**, 147–158
Anoplopoma fimbria: **3**, 146–150
Antarctic dragonfishes phylogeny: **5**, 89–104
Antarctic dragonfishes systematics: **5**, 89–104
Antidepressant effect: **9**, 392–394
Aphanopinae: **9**, 183–192
Arctic char: **7**, 1–9
Arctic grayling: **9**, 220–226
Arctic seas: **1**, 81–123
Arcto-Norwegian cod: **7**, 62–74
Arctozenus rissoi: **4**, 88–96; **6**, 1–19
Artedidraco mirus: **9**, 395–399
Astronesthes chrysophekadion: **2**, 21–39
Astronesthidae: **3**, 121–123
Atlantic halibut: **4**, 150–152
Atlantic salmon: **7**, 52–61

Baikal oilfishes: **1**, 139–145
Baikal omul: **3**, 54–64
Banded flounder biology: **3**, 22–28
Barbourisia rufa: **6**, 140–150
Barbourisiidae: **6**, 140–150
Barents Sea: **4**, 150–152; **5**, 40–54; **6**, 28–44, 45–60; **8**, 245–247
Bathyal region: **4**, 130–134, 135–139
Bathydraconidae: **5**, 89–104
Bathymetric distribution: **9**; 295–304
Behavior of fishes: **1**, 124–138
Behavior of roach: **4**, 121–129
Behavior studies: **2**, 75–98
Beluga sturgeon status: **9**, 59–68
Benthopelagic sculpins: **3**, 54–64
Bering Sea: **3**, 65–77; **4**, 9–17, 18–27; **8**, 24–32
Biochemical estimate: **5**, 136–146
Biological characterization: **4**, 28–45
Biological markers: **1**, 44–51
Biology of cisco: **9**, 368–373
Biology: **6**, 1–19; **8**, 44–52
Biomass: **8**, 44–52
Black Sea: **4**, 1–8, 107–120
Black Sea areas: **9**, 113–122
Bleak: **8**, 173–183
Blicca bjoerkna: **9**, 356–361
Blood serum: **2**, 1–7; **6**, 100–107; **8**, 263–270
Blunt-snouted lenok: **3**, 1–21
Bol'shaya River estuary: **2**, 65–74; **3**, 22–28; **6**, 88–99; **9**, 102–112;
Bottom fishes: **9**, 205–219
Bream: **8**, 1–13
Broken Spur hydrothermal vent field: **9**, 328–332
Bureya River: **9**, 374–378

C. (*Cichlasoma*) *portalegrense*: **7**, 86–107

Calcium: **6**, 108–118
Cambodia: **4**, 81–87
Capelin availability: **7**, 62–74
Capture of large specimens: **4**, 150–152
Carangid species: **8**, 137–149
Carangidae: **3**, 40–53
Carp: **8**, 1–13
Carp sperm: **9**, 305–314
Caspian Sea: **9**, 59–68
Caspian spiny loach: **9**, 362–365
Caspian trout: **2**, 8–20
Catchability coefficient: **2**, 120–127
Cation concentration: **2**, 111–119
Caucasian freshwater gobies: **2**, 139–157
Caucasian gobies differentiation: **1**, 1–20
Champsocephalus gunnari: **7**, 153–157
Channichthyidae: **7**, 153–157
Chars: **8**, 14–23
Chernobyl: **2**, 40–64
Chita Oblast: **9**, 142–153
Chromosomal complement: **9**, 395–399
Chromosome set description: **3**, 132–136
Chukotski Peninsula rivers: **8**, 77–90
Chum salmon: **1**, 146–157; **9**, 40–46; **9**, 171–182
Cichlasoma (*Archocentrus*) *nigrofasciatum*: **7**, 86–107
Cichlasoma octofasciatum: **9**, 135–141
Cichlidae: **7**, 86–107; **9**, 135–141
Clupea harengus: **3**, 132–136
Coastal waters: **4**, 140–144
Cobitidae: **3**, 140–142; **9**, 362–365
Cobitis lebedevi: **3**, 140–142
Cod: **6**, 28–44, 45–60
Coho salmon feeding: **9**, 386–391
Commander Islands: **8**, 248–252
Common carp intestine: **8**, 257–262
Common exploited fishes: **4**, 107–120
Comparative biological characterization: **8**, 77–90
Comparative karyology: **8**, 14–23
Composition changes: **7**, 108–121
Computer identification: **4**, 1–8
Condition of stocks: **9**, 351–355
Contemporary phenotypic diversity: **8**, 207–218
Continental coast: **9**, 40–46
Continental waters: **4**, 56–80
Cooling pond: **2**, 40–64
Copper: **8**, 226–232; **9**, 154–170
Coregonid fish systematics: **5**, 105–135
Coregonid fishes ecology: **5**, 105–135
Coregonidae: **5**, 136–146
Coregonus albula: **9**, 368–373
Coregonus autumnalis migratorius: **3**, 54–64
Coregonus lavaretus pidschian: **5**, 136–146; **8**, 207–218
Coregonus sardinella: **4**, 28–45; **8**, 61–76
Coryphaenoides rupestris: **9**, 123–134
Cottidae: **4**, 140–144; **8**, 245–247; **9**, 89–101
Cottoids: **9**; 295–304
Cottunculidae: **8**, 245–247
Countries adjacent to Russia: **9**, 256–272
Cryopreservation of sperm: **9**, 305–314
Current state of stocks: **9**, 40–46
Cutlass fishes: **2**, 128–138
Cyclops scutifer: **7**, 75–85
Cyclopteridae: **8**, 245–247
Cyclothone: **8**, 53–60
Cyprinidae: **9**, 345–350
Cyprinus carpio: **6**, 108–118; **8**, 1–13, 257–262; **9**, 305–314

Dagi River ichthyocene: **5**, 13–27
Daily dynamics: **8**, 257–262
Daphnia longiremis: **7**, 75–85
Dasyatididae: **5**, 74–88
Deep-sea fishes: **9**, 328–332
Development disturbances: **6**, 72–87
Diagnostic characters of species: **8**, 98–110
Diet: **9**, 113–122
Diet of Baikal omul: **3**, 54–64
Differing levels of pollution: **9**, 113–122
Digestive tract of fishes: **5**, 28–39
Digestive tract protective function: **5**, 28–39
Distinctive features: **8**, 219–225
Distribution: **1**, 59–71; **6**, 140–150; **8**, 44–52 **9**, 154–170
Downstream migration: **9**, 227–248
Dragon stingray: **5**, 74–88

Early development stages: **8**, 111–122
East Atlantic: **9**, 333–337
Eastern Atlantic: **5**, 63–73

Eastern Europe: **4**, 56–80
Eastern redfins: **9**, 351–355
Ecological characteristics: **9**, 8–19
Ecology of fishes: **4**, 145–149
Elasmobranchii: **5**, 63–73; **5**, 74–88; **6**, 151–159
Emmelichthyidae: **6**, 20–27
Encrasicholina punctifer: **3**, 151–156
Engraulididae: **3**, 151–156
Environmental indicators: **5**, 147–158
Environmental influences: **5**, 13–27
Epipelagic fishes: **9**, 193–204
Epipelagic zone: **9**, 273–280
Erythrocyte membranes: **8**, 1–13
Erythrocytes: **2**, 111–119; **6**, 108–118
Esox lucius: **9**, 379–385
Estimate of stocks: **9**, 351–355
Eumicrotremus soldatovi: **8**, 44–52
Exogenous estradiol effect: **3**, 109–120
Exploited bream population: **9**, 69–78
Exploited fish fauna changes: **4**, 56–80
Exploited fish population: **9**, 249–255

Factors determining ontogeny stages: **4**, 107–120
Facultative social strategy: **8**, 167–172
Family Comephoridae: **1**, 139–145
Fecundity: **8**, 245–247
Feeding habits: **2**, 65–74; **5**, 40–54; **6**, 88–99; **8**, 248–252; **9**, 79–88, 89–101, 102–112
Feeding patterns: **9**, 356–361
Feeding strategy: **5**, 40–54
Female parental investment: **9**, 135–141
Finland: **9**, 368–373
First-generation offspring: **2**, 40–64
Fish aggregations: **4**, 97–106
Fish community changes: **5**, 1–12
Fish larvae: **4**, 1–8
Fish schooling behavior: **8**, 167–172
Fish swimming activity: **9**, 392–394
Fishery-regulations theory: **8**, 239–244
Fishes in Russia: **2**, 75–98
Fishlike vertebrates: **1**, 81–123
Flowing water: **8**, 173–183
Flows in Ural River: **1**, 52–58
Fluctuating asymmetry: **9**, 171–182
Food chain: **7**, 122–142
Foraging dynamics: **3**, 65–77
Formal presentation method: **9**, 249–255
Fossil genus *Anenchelum*: **9**, 183–192
Free amino acids: **2**, 8–20
Frequency of occurrence: **8**, 245–247
Freshwater fishes distribution: **8**, 33–43
Functional disturbances: **6**, 72–87
Functional state: **7**, 143–152

Gadidae: **8**, 233–238; **9**, 113–122
Gadus macrocephalus: **4**, 9–17, 18–27; **8**, 248–252
Gadus morhua morhua: **6**, 28–44, 45–60; **7**, 62–74
Gaidropsarus pakhorukovi: **8**, 233–238
Gasterosteidae: **2**, 65–74; **6**, 88–99
Gasterosteus aculeatus: **6**, 88–99
Genetic estimate: **5**, 136–146
Genetic origin identification: **1**, 146–157
Genus: **3**, 40–53
Genus *Aphanopus*: **2**, 128–138
Genus *Brachymystax*: **3**, 1–21
Genus *Cottocomephorus*: **3**, 54–64
Genus *Hexatrygon*: **6**, 151–159
Genus *Lyconus*: **8**, 98–110
Genus *Salmo*: **8**, 198–206
Genus *Salvelinus*: **8**, 77–90
Genus *Thymallus*: **9**, 374–378
Genus *Triglops*: **4**, 140–144
Germ cells: **8**, 123–136
Gobiidae: **1**, 1–20; **2**, 139–157
Gonadal development: **3**, 101–108
Gonadotropic hormones: **2**, 1–7
Gonostomatidae: **8**, 53–60
Gray mullet: **2**, 1–7
Grayling: **9**, 374–378
Growth of roach: **8**, 253–256; **9**, 256–272
Growth: **1**, 59–71; **7**, 62–74
Growth of young: **7**, 75–85
Gustatory properties: **2**, 8–20

Haptenchelys texis: **9**, 328–332
Harpagifer species: **9**, 341–344
Harpagiferidae: **9**, 341–344
Hatching: **6**, 119–129
Hemilepidotus: **9**, 89–101
Hereditary factors: **9**, 20–26

Hexatrygonidae: **6**, 151–159
Himantura draco: **5**, 74–88
Himantura gerrardi: **5**, 74–88
Hippoglossoides platessoides limandoides: **5**, 40–54
Hippoglossus hippoglossus: **4**, 150–152
Hoplostethus vniro: **9**, 333–337
Hormonal regulation features: **8**, 184–197
Hormonal status change: **3**, 86–100; **9**, 315–327
Huso huso: **9**, 59–68
Hypomesus olidus: **5**, 55–62
Hypophthalmichthys molitrix: **2**, 40–64; **3**, 86–100; **9**, 315–327
Hypothalamus: **9**, 315–327

Ichthyocene annual changes: **1**, 21–31
Ichthyocene role: **9**, 273–280
Ichthyocenes: **8**, 150–166
Ichthyofauna: **6**, 61–71; **7**, 108–121
Ichthyofauna changes: **3**, 124–128
Ichthyofauna formation: **8**, 33–43
Ichthyoplankton: **3**, 78–85
Impact of intensive pollution: **3**, 124–128
Indian Ocean: **6**, 130–139; 151–159
Individual fish behavior: **8**, 167–172
Indochina: **8**, 33–43
Influence of abundance: **7**, 75–85
Interannual dynamics: **6**, 28–44
Intraspecific differentiation: **9**, 220–226
Introduction of *Dreissena polymorpha*: **8**, 253–256
Introduction of new species: **5**, 1–12
Investigation of production: **9**, 345–350
Investigation of stocks: **9**, 345–350
Ion transport: **8**, 1–13

Juvenile fish behavior: **2**, 99–110
Juvenile fish body size: **2**, 99–110
Juvenile fish growth: **4**, 46–55
Juvenile fishes: **9**, 227–248
Juvenile pike growth: **9**, 379–385
Juvenile pink salmon: **7**, 143–152
Juvenile sterlet: **3**, 109–120
Juveniles: **8**, 173–183

Kamchatka: **1**, 21–31; **4**, 140–144; **8**, 91–97, 198–206
Kamchatka coastal waters: **9**, 89–101
Kara Sea: **4**, 145–149
Karaginskiy Island: **8**, 24–32
Karakum Canal: **7**, 108–121
Karyological evolution: **9**; 295–304
Karyotype evolution: **8**, 14–23
Karyotypes: **9**; 295–304
Karyotypes description: **9**, 341–344
Khramulya: **9**, 345–350
Kuril–Kamchatka trench: **4**, 130–134

Lake Arakhley: **9**, 142–153
Lake Azabach'ye: **9**, 47–58
Lake Azabach'ye isolate: **8**, 91–97
Lake Baikal: **3**, 54–64; **5**, 136–146; **9**; 295–304
Lake Balkhash: **5**, 1–12
Lake Kenon: **9**, 142–153
Lake Kuril'skoye: **7**, 75–85
Lake Ouluyarvi: **9**, 368–373
Lake Pleshcheyevo: **8**, 253–256
Lake Pskov littoral: **1**, 59–71
Lake Sevan: **9**, 345–350
Lake Vozhe: **7**, 122–142
Lama Lake: **8**, 207–218
Lampetra fluviatilis: **8**, 184–197
Lamprey: **8**, 184–197
Large-fin thornyhead: **8**, 219–225
Lateral nuclei: **9**, 315–327
Lebedev's spined loach distribution: **3**, 140–142
Leucaspius delineatus: **8**, 173–183
Lindberg, Professor G.U.: **3**, 157–158
Liopsetta pinnifasciata: **3**, 22–28
Lipid composition: **8**, 263–270
Lipoproteins: **8**, 263–270
Lofoten: **6**, 45–60
Long-term trends: **9**, 27–39
Lower Bol'shaya River: **2**, 65–74
Lumpenidae: **8**, 245–247
Lycenchelys parini sp. nova: **4**, 130–134
Lycenchelys remissaria sp. nova: **4**, 135–139
Lyconus pinnatus larvae: **8**, 98–110

Mackerel icefish: **7**, 153–157
Macquarie Island: **9**, 341–344

Macrouridae: **9**, 123–134
Macruronus magellanicus: **3**, 29–39
Magellan hake biology: **3**, 29–39
Magnesium: **6**, 108–118
Magnisudis atlantica: **4**, 88–96; **6**, 1–19
Male parental investment: **9**, 135–141
Mallotus villosus villosus: **7**, 62–74
Marine fishery: **1**, 146–157
Marine ichthyofauna: **4**, 81–87
Marine period of life: **7**, 143–152
Maritime Province: **9**, 171–182
Maritime Territory waters: **3**, 143–145
Masu salmon: **9**, 1–7
Maturation rates: **6**, 45–60
Mature ovaries: **8**, 123–136
Mediterranean Sea: **2**, 1–7
Merlangus merlangus euxinus: **9**, 113–122
Merlucciidae: **8**, 98–110
Mesopelagic fishes: **3**, 65–77; **9**, 193–204, 273–280
Micronektonic fishing gear: **2**, 120–127
Microphotolepis schmidti: **6**, 130–139
Mid-Atlantic Ridge: **9**, 328–332
Middle Eocene: **9**, 183–192
Middle Vyatka River: **9**, 8–19
Migration through estuary: **7**, 52–61
Migrations of Atlantic halibut: **4**, 150–152
Migratory chars: **8**, 77–90
Miocene deposits: **8**, 24–32
Monitoring: **6**, 100–107
Moroccan waters: **9**, 338–340
Morphofunctional condition: **9**, 315–327
Morphological characteristics: **9**, 8–19
Morphological differences: **3**, 1–21
Morphological estimate: **5**, 136–146
Morphology of roach: **8**, 253–256
Mortality: **6**, 20–27
Mortality-rate estimates: **9**, 69–78
Moscow River above Moscow: **3**, 129–131
Mouthbrooder: **1**, 72–80
Mugil cephalus: **2**, 1–7
Muscles: **6**, 108–118
Muscles of bream: **2**, 111–119
Museum collections analysis: **1**, 1–20
Myctophidae: **8**, 24–32; **9**, 79–88

Natural-mortality estimates: **7**, 153-157
Natural reproduction: **9**, 281-294
Near-bottom fishes: **9**, 205–219
Neogobius platyrostris constructor: **1**, 1–20
Neogobius rhodioni: **2**, 139–157
Neonesthes capensis larva: **3**, 121–123
Neurosecretory cells: **9**, 315–327
New fishing regulations: **9**, 281–294
New form: **8**, 198–206
New records: **2**, 128–138; **3**, 146–150; **9**, 193–204
New species: **2**, 128–138; **2**, 139–157; **8**, 24–32, 233–238; **9**, 183–192, 333–337
New species description: **2**, 21–39
Nickel: **9**, 154–170
Nine-spined stickleback: **2**, 65–74
Norilsk-Pyasina lakes: **8**, 207–218
Norilsk-Pyasinskiy water system: **3**, 124–128
Norilsk–Pyasin water system: **8**, 61–76
North American distribution: **5**, 55–62
North Asia: **4**, 56–80
North Atlantic: **9**, 123–134
North Caspian basin: **9**, 362–365
Northeast Pacific Ocean: **3**, 121–123
Northeast Russia: **9**, 220–226
Northeast Sakhalin: **5**, 13–27
Northern Caucasus: **9**, 183–192
Northwest Pacific: **6**, 1–19; **9**, 386–391
Northwest Pacific Ocean: **4**, 88–96
Northwest Sakhalin: **1**, 146–157
Norwegian Sea: **5**, 40–54
Notal waters: **8**, 111–122
Notoscopelus kroeyerii: **9**, 79–88
Notothenioidei: **5**, 89–104; **9**, 341–344, 395–399
Nuclear power plant: **2**, 40–64
Number of vertebrae: **9**, 20–26

Oncorhynchus gorbuscha: **7**, 42–51; 143–152
Oncorhynchus keta: **1**, 146–157; **9**, 40–46; 171–182
Oncorhynchus kisutch: **9**, 386–391
Oncorhynchus masou: **9**, 1–7
Oncorhynchus mykiss: **3**, 137–139
Oncorhynchus nerka: **7**, 75–85; **8**, 91–97; **9**, 47–58

Ontogeny: **3**, 1–21
Ontogeny in bream: **3**, 101–108
Oogenesis: **1**, 72–80; **6**, 119–129; **8**, 137–149
Oreochromis mossambicus: **1**, 72–80; **6**, 119–129;
Osmeridae: **5**, 55–62
Osmerus eperlanus: **9**, 368–373
Osmoregulatory system: **7**, 143–152
Osteology: **6**, 140–150
Ovarian development: **3**, 109–120
Ovarian follicles: **3**, 86–100

Pachycara thermophilus: **9**, 328–332
Pacific basin trout: **8**, 198–206
Pacific cod: **4**, 9–17; **8**, 248–252
Pacific cod abundance: **4**, 18–27
Pacific cod behavior: **4**, 18–27
Pacific cod biology: **4**, 9–17
Pacific cod distribution: **4**, 18–27
Pacific cod exploitation: **4**, 9–17
Pacific Ocean: **8**, 53–60
Pacific Ocean shores of Japan: **4**, 135–139
Pacific waters: **1**, 21–31; **9**, 193–204
Parasites: **1**, 44–51
Pelecus cultratus: **3**, 129–131
Perciformes: **4**, 130–134; 135–139; **5**, 89–104
Peru: **9**, 227–248
Peter the Great Bay: **9**, 351–355
pH fluctuation influence: **4**, 46–55
Philippine Sea: **3**, 151–156
Physicochemical parameters: **6**, 100–107
Physiological condition: **3**, 137–139
Phytophilous fish species: **1**, 59–71
Pink salmon: **7**, 42–51
Plagiogeneion rubiginosum: **6**, 20–27
Plasma: **2**, 111–119; **6**, 108–118
Pleuronectidae: **3**, 22–28
Pogonophryne dolichobranchiata: **9**, 395–399
Pollution from human activity: **3**, 124–128
Pond rearing: **8**, 257–262
Pond smelt: **5**, 55–62
Population differences: **9**, 1–7
Population differentiation: **5**, 136–146
Population dynamics: **7**, 10–41; **9**, 47–58
Population structure: **7**, 42–51
Potassium: **6**, 108–118
Predatory fishes: **9**, 102–112
Premigration: **7**, 143–152
Preoptic nuclei: **9**, 315–327
Prespawning behavior differences: **7**, 86–107
Prespawning period: **3**, 86–100; **9**, 315–327
Primor'ye: **9**, 1–7
Ptilichthys goodei: **3**, 143–145
Pungitius pungitius: **2**, 65–74

Quaternary: **4**, 56–80
Quillfish: **3**, 143–145

Radiobiological analysis: **2**, 40–64
Rainbow trout: **3**, 137–139
Rainbow trout physiology: **8**, 226–232
Raja clavata: **5**, 63–73
Rajidae: **5**, 63–73
Rare deepwater fishes: **8**, 111–122
Rare mesopelagic fishes: **4**, 88–96; **6**, 1–19
Rare species: **6**, 130–139
Ration: **6**, 28–44
Reaction to hydrostatic pressure change: **8**, 173–183
Recruitment: **7**, 10–41; **8**, 123–136; **9**, 47–58
Recruitment theory: **7**, 10–41
Reduced temperature effect: **1**, 72–80
Reduced temperature effect: **6**, 119–129
Reproduction function: **8**, 184–197
Reproduction: **1**, 59–71
Reproductive cycle: **9**, 123–134
Reproductive period: **5**, 13–27
Reproductive system: **6**, 72–87
Reproductive system development: **2**, 40–64
Reproductive variability of populations: **8**, 239–244
Reservoirs: **9**, 356–361
Retinal structure: **1**, 139–145
Review: **5**, 105–135
Rio Grande Rise: **8**, 233–238; **9**, 205–219
Riverine period of life: **7**, 143–152
Roach: **1**, 44–51; **9**, 20–26
Roach ecomorphs: **1**, 44–51
Rockling: **8**, 233–238
Role of heterochrony: **3**, 1–21
Roundnose grenadier: **9**, 123–134
Running water: **4**, 121–129

Russia: **9**, 256–272
Russian studies: **1**, 124–138
Russian sturgeon: **6**, 100–107; **8**, 123–136
Rutilus rutilus: **1**, 44–51; **4**, 121–129; **8**, 253–256; **9**, 20–26, 256–272
Rybinsk Reservoir: **1**, 44–51; **2**, 111–119; **3**, 101–108

Sabanejewia caspia: **9**, 362–365
Saberfish: **3**, 129–131
Sabkhet el Bardawîl: **2**, 1–7
Sablefish: **3**, 146–150
Sakhalin–Kuril region: **1**, 32–43
Salmo gairdneri: **8**, 226–232
Salmo salar: **7**, 52–61
Salmo trutta caspius: **2**, 8–20
Salmonidae: **3**, 1–21
Salvelinus: **7**, 1–9
Scabbardfish: **9**, 183–192
Schooling interactions: **2**, 99–110
Scopelosaurus harryi: **4**, 88–96; **6**, 1–19
Scorpaenidae: **8**, 219–225
Scotia Sea: **9**, 395–399
Sculpins: **4**, 140–144
Sculpins abundance: **3**, 54–64
Sculpins biomass: **3**, 54–64
Sea of Japan: **9**, 366–367
Sea of Okhotsk: **3**, 146–150; **8**, 44–52; **9**, 40–46, 273–280, 386–391
Seasonal changes: **2**, 1–7
Seasonal dynamics: **6**, 28–44
Seaward migration: **7**, 52–61
Sebastolobus macrochir: **8**, 219–225
Second record: **6**, 151–159
Secretory cells: **8**, 123–136
Sensitivity to hormones: **3**, 86–100
Sensory systems of fishes: **1**, 124–138
Sensory-system studies: **2**, 75–98
Sex differentiation: **3**, 101–108
Sex steroid hormones: **2**, 1–7
Sex structure: **4**, 140–144
Sexual cycle final period: **8**, 184–197
Sharp-snouted lenok: **3**, 1–21
Siberian cisco: **4**, 28–45; **8**, 61–76
Silver carp females: **9**, 315–327
Silver carp: **2**, 40–64
Silver-carp females: **3**, 86–100
Sixgill stingray: **6**, 151–159
Size: **4**, 140–144
Size–age composition: **9**, 113–122
Size composition: **8**, 219–225
Slickheads: **6**, 130–139
Smelt: **9**, 368–373
Smolt age: **8**, 91–97
Sockeye salmon: **8**, 91–97; **9**, 47–58
Sodium: **6**, 108–118
Soldatov's spiny lumpsucker: **8**, 44–52
South Georgia Island area: **7**, 153–157
South Orkney Islands: **9**, 341–344
Southeastern Sakalin: **8**, 219–225
Southern Kuril Islands: **9**, 193–204
Southern Madagascar Ridge: **5**, 63–73
Southern Sakhalin: **7**, 42–51; **9**, 1–7
Southwest Atlantic: **3**, 29–39; **8**, 233–238
Southwest Atlantic Ocean: **9**, 205–219
Southwest Bering Sea: **9**, 27–39
Southwestern Pacific: **8**, 98–110, 111–122
Spatial distribution: **6**, 1–19; **8**, 219–225
Spatial distribution of fish: **9**, 142–153
Spatial structure: **8**, 91–97
Spawning stock: **4**, 28–45
Species key: **3**, 40–53
Species trophic status: **8**, 150–166
Spined form: **9**, 338–340
Stages of ontogeny: **4**, 107–120
Stellate sturgeon: **9**, 281–294
Stellate sturgeon recruitment: **1**, 52–58
Sterlet: **9**, 8–19
Stizostedion lucioperca: **7**, 122–142
Stocks: **9**, 47–58
Structure of fish aggregations: **4**, 97–106
Sturgeons: **6**, 72–87; **8**, 263–270
Subarctic waters: **9**, 154–170
Subgenus *Stomianodon*: **2**, 21–39
Subisolates: **9**, 47–58
Sublethal concentrations: **8**, 226–232
Suspended particles' effect: **4**, 121–129
Synaphobranchidae: **9**, 328–332
Syr Darya basin: **6**, 61–71
Syrian surface waters: **3**, 78–85
Systematic: **3**, 40–53
Systematics: **2**, 139–157

Tactostoma macropus: **4**, 88–96; **6**, 1–19
Taimyr Peninsula: **7**, 1–9; **8**, 207–218
Taxon: **3**, 40–53
Taxonomic descriptions: **4**, 88–96
Teleostei: **3**, 40–53
Territorial interactions: **2**, 99–110
Thermophilic fish species: **9**, 366–367
Thornback skate: **5**, 63–73
Three-spined stickleback: **6**, 88–99
Thymallus arcticus: **9**, 220–226
Tilapia: **6**, 119–129
Tissues of fishes: **9**, 154–170
Total carbohydrase activity: **8**, 257–262
Trachichthyidae: **9**, 333–337
Trachurus: **3**, 40–53
Trichiuridae: **2**, 128–138; **9**, 183–192
Trophic relationships: **9**, 79–88
Trophic status determination methods: **8**, 150–166
Trophic system: **9**, 89–101
Turkmenistan: **7**, 108–121

Unstable recruitment level: **9**, 69–78
Upper epipelagic zone: **1**, 21–31; **1**, 32–43
Ural River: **1**, 52–58

Varicorhinus capoeta sevangi: **9**, 345–350
Verkhovka: **8**, 173–183
Vertical distribution: **8**, 53–60
Volga River: **9**, 281–294

Walters Shoal: **5**, 63–73
Water temperature: **7**, 62–74
Waters adjacent to Arctic seas: **1**, 81–123
West Bering Sea: **1**, 21–31
West Indian Ocean: **8**, 137–149
West Indian submarine ridge: **6**, 20–27
West Kamchatka: **2**, 65–74; **6**, 88–99; **9**, 102–112
Western Indian Ocean: **5**, 74–88
Western Kamchatka: **3**, 22–28
White Sea: **3**, 132–136
Whitefish: **5**, 136–146
Whiting: **9**, 113–122
Winter nektonic communities: **9**, 27–39

Xenobiotic influence: **8**, 1–13

Yenisey River: **6**, 72–87
Young white bream: **9**, 356–361

Zander: **7**, 122–142
Zeidae: **9**, 338–340
Zeus faber: **9**, 338–340
Zinc: **9**, 154–170
Zoarcidae: **4**, 130–134, 135–139; **8**, 245–247; **9**, 328–332